



3-A-Day™ dairy keeps dental caries away



Despite the widespread use of fluoride in an effort to reduce tooth decay, dental caries remain one of the most common childhood diseases. Although not life threatening, caries cause pain and discomfort, resulting in reduced food intake and therefore affecting quality of life. Dental caries is a multifactorial disease that is affected by physical and biological factors (maternal characteristics, morphology and composition of teeth, cariogenic bacteria and fluoride exposure), lifestyle and behavioural factors (oral hygiene practices and dietary habits), epigenetics and social status.¹ Dairy products offer many health benefits beyond the well-known contribution to building strong bones, specifically with regard to preventing dental caries and periodontitis. Milk not only has carioprotective characteristics due to the buffering activity of milk protein, but also contains various bioactive peptides, calcium, casein and lipids, which have proven to be important in the maintenance of dental health because of their associated cariostatic properties.¹⁻⁴

The facts

There is substantial evidence for the role of milk products in promoting dental health.

- Several studies support the association between the consumption of milk or milk products and the occurrence of fewer caries due to the presence of bioactive compounds.^{1,2,5-7}
- Milk consumption does not increase plaque acidity, which makes it essentially a 'tooth-friendly' food.^{1,2,8,9}
- Cheese significantly increases the concentration of calcium in saliva and plaque, which confers a cariostatic effect. Eating cheese (especially aged cheese) before or after sugary foods prevents a drop in the pH of plaque and therefore confers an enamel-protecting effect.^{1,5,6,10}
- A low dietary intake of calcium may lead to the development of periodontal disease, whereas increased calcium and vitamin D intake seems to facilitate a protective effect that prevents tooth loss.^{1,8,9,11,12}

- Studies have shown that adding milk, or the milk ions calcium and phosphate, to food not only diminishes the extent of tooth demineralisation but may also promote remineralisation.^{1,3,6,7,13}
- A high consumption of yoghurt may be associated with a lower prevalence of dental caries in young children.^{1,6,7,9,11,14,15}

What is dental decay?

Dental decay refers to the development of dental caries and the subsequently compromised underlying periodontal tissue. Tooth enamel is a polymeric substance that consists of crystalline calcium phosphate embedded in a protein matrix. Dental caries form when calcium and phosphate of the tooth enamel dissolves owing to the action of acid in the mouth (acidic demineralisation). Demineralisation can occur directly (when acidic food is consumed) or indirectly (by fermentation products produced by oral bacteria growing on residual food particles between teeth or adhering to plaque).^{2,3}

Benefits of dairy products for dental health

Milk and milk products may have extra health benefits that could apply to a broader dental perspective owing to calcium, phosphate and various protein structures found in dairy. Several studies substantiate the beneficial effect of dairy on dental health. Evidence shows that milk or dairy products can reduce the demineralisation of enamel, promote its remineralisation and prevent the adhesion of *Streptococcus mutans* to the surface of teeth.^{15,16} The concentration of calcium in dental plaque and that of calcium and phosphate ions in the saliva affect the balance between demineralisation and remineralisation of enamel. Calcium, phosphate, casein, proteins and fats in dairy are considered to be anticariogenic, therefore offering further dental protection.

A study among 12 764 American adults (older than 18 years) showed an inverse association between the intake of dairy and the prevalence of periodontitis. Individuals who consumed milk products more than four times a week were 20% less likely to present with periodontitis than those who consumed dairy only once a week or less.^{1,8} Also, higher maternal intake of total dairy products, yoghurt and calcium during pregnancy was found to be associated with a reduced risk of dental caries in children.^{1,9,11,17} Higher intakes of calcium and dairy servings are also reported to be associated with lower plaque scores, especially when the vitamin D intake exceeds 6.8 µg/day.⁸

A long-term study investigating the development of caries among children and adolescents, showed that 72% of the participants with a dairy intake above the mean had no caries, compared with 65% who reported below-average dairy intake.¹⁸ The intake of dairy and milk, and also of dairy-derived nutrients such as calcium, whey and casein, was found to be inversely associated with childhood and adolescent caries. Dairy intake in early childhood was further observed to be a

potential predictor of future risk of caries, based on the number of decayed, missing or filled surfaces of teeth.

Cheese

When hydrolysed for long periods, casein generates bioactive peptides called casein phosphopeptides (CPPs). CPPs are sections of casein with three phosphoserines followed by two glutamic acid residues, capable of binding the calcium phosphate present in milk. CPPs were first discovered in cheese where they exhibited anticariogenic effects in humans and animals. Consumption of milk products in general showed to decrease dental caries, but the result with cheese was more pronounced. This is due to the sticky nature of cheese, which allowed it to adhere to the teeth. When in close proximity to the tooth enamel, CPPs act as a mineral reserve, which replenishes the tooth's calcium and phosphate lost owing to acid erosion from food or oral bacteria. CPPs have the added effect of reducing the adhesion of cariogenic bacteria, thereby further decreasing the development of dental caries.¹⁰

Chewing hard cheese, particularly Cheddar cheese, has been found to prevent enamel demineralisation not only by stimulating salivary flow, which buffers the pH of dental plaque, but also by increasing the calcium and phosphorus concentrations in dental plaque, which favours remineralisation.¹² A prospective study from Japan also found that higher maternal intake of cheese during pregnancy was associated with a decreased risk of dental caries in children.¹

Yoghurt

Regular yoghurt intake (≥ 4 times/week) was found to be significantly associated with lower prevalence of caries compared with intakes of less than once a week, showing a clear dose–response relationship.^{1,9} A study investigating the relationship between the intake of various milk products and periodontitis concluded that lactic-acid foods such as yoghurt may be beneficial in preventing periodontal disease.¹ Similar results were obtained in a cross-sectional study among Italian toddlers (3–5 years old), with the consumption of yoghurt shown to be significantly inversely associated with rampant early childhood dental decay.^{9,19}

Flavoured milk products

It is reasonable to assume that adding sucrose to milk will increase cariogenicity. However, the concentration at which the added sucrose overcomes the carioprotective properties of plain milk needs to be considered. The question is complicated by cocoa's protective properties in chocolate-flavoured milk.²⁰ Recent findings indicate that the cariogenic load of flavoured milk products is negligible to low.^{13,21} This is supported by the observation that children who consumed conventional chocolate milk (which contains cocoa and sugar) regularly over a two-year period had a slight but non-significant increase in caries compared to those who drank plain, white milk.^{3,21}

Therefore, although the sugar content of flavoured milk has raised health concerns, sweetened dairy is regarded to be a preferable alternative to similarly sweetened beverages such as cordials, soft drinks and citrus juices. Regular consumption of the latter two examples promotes dental erosion through direct acidic demineralisation rather than through increased bacterial activity because of a high sugar content. The high concentration of calcium and phosphate in dairy buffers the acidic action associated with bacterial metabolism. Sweetened dairy products can also promote milk consumption among both adults and children, which delivers essential protective nutrients such as casein, calcium and phosphorus.^{13,21}

To understand the effect of sugar on cariogenicity, the effect of

10% additional sucrose or fructose in milk was studied.¹³ There was little difference in cariogenicity between these two sweetened milk products and although both were more cariogenic than plain, white milk, they were still less cariogenic than 10% sucrose in water. The experiment also showed that 4% lactose in water had a very low cariogenicity, which suggests that the natural sugar content of milk is unlikely to promote the development of caries. This observation is supported by a study which showed that despite the addition of 2% sucrose to milk enhancing its cariogenic activity, such sweetened milk is significantly less cariogenic than similar water–sucrose solutions.¹³

Findings from a number of studies therefore suggest that the benefits of moderate consumption of flavoured milk outweigh possible negative effects, particularly if efforts are made to find a balance between trying to increase milk consumption and control sugar intake.²¹

Potential mechanisms of protection

In addition to the nutritional benefits of dairy, an increasing body of evidence demonstrates a bioactive function of dairy components in preventing dental caries among adults.⁷ The non-cariogenic and cariostatic properties of milk appear to be related to several factors.

The carioprotective effect of dairy is thought to be associated with a number of mechanisms that involve more than the presence of calcium (although these are only partially understood). Research has shown that the removal of casein, fat or lactose does not affect the protective remineralisation potential of milk. Although the removal of free calcium and phosphate showed some effect, water-soluble proteose peptones, which were found to be powerful protective factors, were still present.⁷

Phosphopeptides

Milk contains two major protein groups, namely insoluble caseins and soluble whey proteins, distinguished according to their solubility in unheated milk (pH 4.6; 20 °C). Both groups have specific physiochemical and biological properties.² Bioactive proteins and peptides are embedded in primary sequences in casein and whey.^{2,8} Milk-derived bioactive sequences are involved in a variety of regulatory activities, such as modulating digestive and gastrointestinal functions, regulating haemodynamics (hypertension and gastric blood flow), conferring anticariogenic and analgesic properties, synthesis of growth factors and immunoregulation. Caseins, which account for 80% of milk proteins, contain several bioactive peptides. In general, bioactive peptides with anticariogenic activity prevent dental lesions through several mechanisms, including bacterial inhibition, competitive exclusion to enamel binding sites, improved buffering capacity in the pellicle surrounding teeth, reduced enamel demineralisation and enhanced enamel remineralisation.^{5,10}

The adherence of oral bacteria to saliva-coated hydroxylapatite of tooth enamel has been found to be inhibited specifically by three milk-derived compounds, namely CPP, sodium caseinate and glycomacropeptide.^{4,22} Research showed that a paste of CPP and amorphous calcium phosphate (CPP–ACP), found in higher concentrations in yoghurt than in equal-weight portions of cheese and milk, can bind and stabilise calcium and phosphate in solution. CPP–ACP also influences the demineralisation/remineralisation process of dental enamel by binding to both dental plaque and tooth enamel.^{2,3,5,10,11}

Lactose

Lactose cariogenicity has been intensely debated for many years. Much of this debate stems from evidence that organic acid metabolites produced from lactose fermentation by the

oral microflora have a deleterious effect on tooth enamel. Although lactose comprises 80% of the carbohydrate content of milk – sufficient to render milk cariogenic – evidence shows that lactose is the least cariogenic of the common dietary sugars. Also, the co-presence of bioactive components in dairy has been found to provide a buffering action against the possible cariogenicity of lactose.^{13,23} The high concentrations of calcium and phosphorus in milk help to prevent dissolution of enamel (which consists mainly of calcium and phosphate), while other components of milk may reduce the ability of plaque microflora to adhere to enamel and produce acids. The lactose content in fermented dairy products (e.g. yoghurt) was found to be substantially reduced, supporting low cariogenic capability. Apart from the reduced lactose content in fermented milk products, lactic acid also lowers the pH, which inhibits growth of many pathogenic organisms.^{1,6,14}

Probiotics

The use of beneficial bacteria in the field of dentistry has only recently gained interest. Investigation of the effects of probiotic bacteria on oral health has become an important research subject. Although these studies are still in the early stages, results show that probiotic bacteria are effective against tooth caries, periodontal disease, oral mucosal lesions and oral malodour.²⁴ Probiotics are defined as live microorganisms that, when administered in adequate amounts, confer a health benefit on the host through balancing the microbial flora. Probiotics may have beneficial effects in the mouth, possibly by changing the level of pathogenic bacteria present in the oral cavity and dental plaque. This is likely due to the displacement of *S. mutans* or other cariogenic microorganisms from their binding sites on dental surfaces or in oral biofilms. Studies have shown that probiotics in milk products such as yoghurt reduce *S. mutans* counts, possibly by modifying the composition of the salivary film, which prevents bacterial adhesion. This finding is supported by a study which showed *S. mutans* counts decreasing significantly with consumption of probiotic yoghurt compared with plain yoghurt.^{3,6,14} Consumption of products that contain probiotics is currently considered an appropriate method in preventing dental caries.⁵

An interesting recent study showed research focus shifting from the caries-promoting bacterium *S. mutans* only to a more complex acid-tolerant consortium of oral bacteria implicated in disease. Molecular microbiological methods have shown that even with a sugar-rich diet, a much broader spectrum of acidogenic microbes are found in the dental biofilm. Prevention of tooth decay by targeting solely *S. mutans* or by mainly limiting sugar intake would not be sufficient to protect against disease progression. Bacteriotherapy has emerged as an alternative preventive measure towards a healthier oral microbial community. Results have shown *in vitro* that caries pathogens could be effectively inhibited by oral lactobacilli, with *Lactobacillus* SD1-SD6 having the strongest inhibitory effect against *S. mutans* and *S. sobrinus*.²⁵ Indeed, in different carriers, such as milk and cheese, *Lactobacillus* spp. have shown to have a caries-reducing effect in children.^{9,19} The caries-preventive effect of *Lactobacillus* could be due to improved salivary buffering capacity and reducing the counts of more than one type of bacterium.

Lactoferrin

Lactoferrin, an iron-binding protein found in cow's milk also inhibits the aggregation and surface adherence of Gram-negative bacteria such as *S. mutans*, a major bacterial contributor to development of dental caries.² The inhibitory effect of bovine milk on saliva aggregation of *S. mutans* is

ascribed to the interference of lactoferrin with a streptococcal surface protein antigen.^{7,4}

Vitamin D

Children and adolescents are particularly vulnerable to the clinical manifestations of insufficient vitamin D owing to these developmental stages being critical periods for bone and tooth formation. The function of vitamin D in tooth development implies that impaired tooth composition is more prevalent in subjects with vitamin D deficiency, but the association may be overlooked as clinical manifestations appear only after a considerable delay. Data from a cross-sectional, nationally representative sample of Canadian children suggest that there is an association between caries and lower serum vitamin D. Improving children's vitamin D status with milk-based vitamin D supplementation may be an additional preventive consideration to lower the risk for caries.¹²

Conclusion

There is consistent scientific evidence that milk and milk products are beneficial to dental health. Both the low cariogenic potential of milk and its carioprotective role have been demonstrated. The protective action appears to be facilitated by several properties of milk, namely: (1) low cariogenic potential of lactose; (2) casein (and possibly also milk fats) being carioprotective; (3) the presence of bioactive peptides, and (4) the protective and buffering role of calcium and phosphorus. The addition of probiotics to milk may also benefit oral health, although evidence is limited at present. The benefits of dairy for dental health should be advocated as a complement to proper oral hygiene to help reduce the incidence of dental caries.





References

1. Tanaka K, Miyake Y, Sasaki S, Hirota Y. Dairy products and calcium intake during pregnancy and dental caries in children. *Nutr J*. 2012; 11(33):1-8.
2. Aimutis WR. Bioactive properties of milk proteins with particular focus on anticariogenesis. *J Nutr*. 2004; 134(4):989S-995S.
3. Al-Dajani M, Limeback H. Emerging science in the dietary control and prevention of dental caries. *J Calif Dent Assoc*. 2012; 40(10):34-38.
4. Adegboye AR, Rodrigues Tomaz M, Jeavons CA, Heitmann BL. Intake of dairy foods and oral health: Review of epidemiological data. *Tandlægebladet (Danish Dental Journal)*. 2017; 120(12):884-891.
5. Tunick MH, Van Hekken DL. Dairy products and health: Recent insights. *J Agric Food Chem*. 2015; 63(43):9381-9388.
6. Ravishankar TL, Yadav V, Tangade PS, Tirth A, Chaitra TR. Effect of consuming different dairy products on calcium, phosphorus and pH levels of human dental plaque: A comparative study. *Eur Arch Paediatr Dent*. 2012; 13(3):144-148.
7. Johansson I. Milk and dairy products: Possible effects on dental health. *Scand J Nutr*. 2002; 46(3):119-122.
8. Adegboye ARA, Christensen LB, Holm-Pedersen P, Avlund K, Boucher BJ, Heitmann BL. Intakes of calcium, vitamin D, and dairy servings and dental plaque in older Danish adults. *Nutr J*. 2013; 12:61.
9. Tanaka K, Miyake Y, Sasaki S. Intake of dairy products and the prevalence of dental caries in young children. *J Dent*. 2010; 38(7):579-583.
10. Noren NE. Creation of a sticky coating of dairy proteins containing bioactive peptides to reduce dental caries. MSc dissertation. Winnipeg: University of Manitoba, 2018. <https://mspace.lib.umanitoba.ca/handle/1993/31027>
11. Dror DK, Allen LH. Dairy product intake in children and adolescents in developed countries: Trends, nutritional contribution, and a review of association with health outcomes. *Nutr Rev*. 2013; 72(2):68-81.
12. Gyll J, Ridell K, Öhlund I, Karlstrand Åkeson P, Johansson I, Holgerson PL. Vitamin D status and dental caries in healthy Swedish children. *Nutr J*. 2018; 17(11):1-10.
13. Rugg-Gunn A, Woodward M. Milk and oral health. 2011. Available from <http://www.borrowfoundation.org/assets/uploads/milk-and-oral-health.pdf>.
14. Poureslami H, Pishbin L, Eslaminejad Z, Moqadam FJ, Farokhi MR. The effects of a dairy probiotic product, Espar, on salivary calcium and *Mutans* streptococci. *J Dent Res Dent Clin Dent Prospects*. 2013; 7(3):147-151.
15. Lin H, Lin J, Hu S, Kuo H, Yang Y. Association of dietary calcium, phosphorus, and magnesium intake with caries status among schoolchildren. *Kaohsiung J Med Sci*. 2014; 30(4):206-212.
16. Hujoel PP, Lingström P. Nutrition, dental caries and periodontal disease: A narrative review. *J Clin Periodontol*. 2017; 44(18): S79-S84.
17. Fernando S, Speicher DJ, Bakr MM, et al. Protocol for assessing maternal, environmental and epigenetic risk factors for dental caries in children. *BMC Oral Health*. 2015; 29(15):167.
18. Lempert SM, Christensen LB, Froberg K, Raymond K, Heitmann BL. Association between dairy intake and caries among children and adolescents: Results from the Danish EYHS follow-up study. *Caries Res*. 2015; 49(3):251-258.
19. Biria M, Amini M, Babaei M, Ahmady AE, Lando HA, Amirabadi GE. Dietary habits and caries experience among 9-year-old school children. *JMED Res*. 2015; 2015:336945.
20. Latif R. Chocolate/cocoa and human health: A review. *Neth J Med*. 2013; 71(2):63-68.
21. Li XE, Drake M. Sensory perception, nutritional role, and challenges of flavored milk for children and adults. *J Food Sci*. 2015; 80(4):R665-R670.
22. Agarwal R, Singh C, Yeluri R, Chaudhry K. Prevention of dental caries – measures beyond fluoride. *J Oral Hyg Health*. 2014; 2(1):122.
23. Aimutis WR. Lactose cariogenicity with an emphasis on childhood dental caries. *Int Dairy J*. 2012; 22(2):152-158.
24. Gungor OE, Kirzioglu Z, Kivanc M. Probiotics: Can they be used to improve oral health? *Benef Microbes*. 2015; 6(5):647-656.
25. Meurman JH, Stamatova IV. Probiotics: Evidence of oral health implications, *Folia Med (Plovdiv)*. 2018; 60(1):21-29.